# POSTOPERATIVE INSTRUCTIONS SHOULDER ARTHROSCOPY WITH SUBACROMIAL DECOMPRESSION

#### ❖ SLING/MOVEMENT

- You may wear the sling as tolerated for comfort. Commonly, patients remove the sling within 3 days. You may wear the sling at night as well if it is more comfortable to do so.
- You may move your shoulder in any direction that is not painful. Do not force the motion though because this will likely cause pain.

## **❖** ICE

An ice machine will be provided to you prior to your surgery. This will help decrease swelling and pain after your surgery. Use the ice machine as much as possible when you get home at intervals of 30 minutes on and off. You should keep the ice machine for approximately two weeks. Do not use the ice machine while you are sleeping.

#### MEDICATIONS

• If you were given a nerve block for anesthesia, it will wear off over 18-24 hours. During this time you will have little to no feeling in the body part where you had surgery (i.e. arm).

To control your pain during this transition while the nerve block is wearing off, you are to eat first and then begin taking the **pain medication** (e.g. Vicodin, Percocet, etc) immediately when you get home from surgery. This will prevent you from having severe pain. Take the pain medication every 4 hours until you go to bed.

• The day after surgery you can take 600 mg of Ibuprofen (Advil/Motrin) every 6 hours to help with inflammation and pain. This medication will help cut down the use of narcotic-based pain medication. However, if you still have pain after taking the Ibuprofen, continue taking the pain medication every 4-6 hours as needed.

## ❖ DRESSING/BANDAGES

- Your shoulder dressing is waterproof. You may shower the next day after surgery. Two days after surgery you may remove the entire dressing. Gently wash the wounds with soap and warm water and towel dry. You may then cover the incisions with band-aids until your post op visit.
- Do not take a bath or submerge your shoulder in water until your incisions are checked at your first post-op visit.

### **❖** APPOINTMENT

• Please call the office prior to, or immediately following, your surgery in order to schedule a postoperative appointment. This should be scheduled 7-10 days after surgery. At that visit your stitches will be removed and you will be given a prescription to begin physical therapy.

If you have any questions or concerns, please contact our office at (310) 860-3048 or email us at surgery@snibbeorthopedics.com.